

Coherence of the Antarctic ice core data from the Vostok station and EPICA

Mokhov I.I.^{1,2}, Nyrov A.O.²

¹A.M. Obukhov Institute of Atmospheric Physics RAS

²Lomonosov Moscow State University

mokhov@ifaran.ru

Consistency of the Antarctic ice core data for temperature and CO₂, CH₄ and dust (D) concentrations in the atmosphere from the Russian Vostok station [1] and EPICA Dome C (EDC) [2] for the past more than 400 ka is estimated with the use of cross-wavelet analysis and linear regressions [3] (see also [4-7]).




Figure 1. Coherence of variations of temperature (a), CO₂ (b), CH₄ (c) and dust (d) concentrations in the atmosphere from EDC and Vostok ice core records. Ordinates – periods in kyr.

Figure 1 shows coherence of variations of temperature T (a), q_{CO_2} (b), q_{CH_4} (c) and q_D (d) concentrations in the atmosphere from EDC and Vostok ice core records. The obtained results show a more significant agreement between the paleoreconstructions of EPICA and Vostok for the longest-term variations with a period of about 100 kyr or more. For shorter periods, the consistency of different data is best within the past 200 kyr.

Table 1. Coefficients of linear regressions of EPICA data (T , q_{CO_2} , q_{CH_4} , q_D) on corresponding Vostok data for different 100 ka time intervals BP (I, II, III, IV and I+II+III+IV): k – regression coefficients, r – correlation coefficients.

		I+II+III+IV 0-400 kyr BP	I 0-100 kyr BP	II 100-200 kyr BP	III 200-300 kyr BP	IV 300-400 kyr BP
k (r)	T	1.32 (0.72)	1.49 (0.90)	1.73 (0.94)	0.78 (0.45)	0.79 (0.41)
	CO ₂	0.83 (0.86)	0.98 (0.94)	0.93 (0.97)	0.52 (0.54)	0.74 (0.77)
	CH ₄	0.73 (0.71)	0.77 (0.74)	0.76 (0.86)	0.42 (0.40)	0.54 (0.51)
	D	0.54 (0.51)	0.87 (0.79)	0.76 (0.74)	0.07 (0.06)	0.46 (0.38)

Table 1 shows coefficients of linear regressions of EPICA data (T , q_{CO_2} , q_{CH_4} , q_D) on corresponding Vostok data for different 100 ka time intervals BP (I, II, III, IV and I+II+III+IV): k – regression coefficients, r – correlation coefficients. According to Table 1 the best agreement of different data is displayed for the past 200 kyr (I,II).




Figure 2. Coefficients of linear regressions of EPICA data on corresponding Vostok data for moving 100 ka time intervals in dependence on time (ka) BP: (a) T , (b) q_{CO_2} , (c) q_{CH_4} , (d) q_D . Black lines - coefficients of regression, green lines – coefficients of correlation.

Figure 2 shows coefficients of linear regressions of EPICA data on corresponding Vostok data for moving 100 ka time intervals in dependence on time (ka) BP: (a) T , (b) q_{CO_2} , (c) q_{CH_4} , (d) q_D . According to Fig. 2 the best consistency of different data is exhibited within the past 200 kyr, especially for q_{CO_2} .

As a whole, the results obtained show the best agreement between EPICA and Vostok data within the past 200 kyr.

References

1. Petit J.R., Jouzel J., Raynaud D. et al., 1999: Climate and atmospheric history of the past 420000 years from the Vostok ice core, Antarctica. *Nature*, 399, 429-436.
2. EPICA community members, 2004: Eight glacial cycles from an Antarctic ice core. *Nature*, 429, 623-628.
3. Grinsted A., Moore J.C., Jevrejeva S., 2004: Application of the cross wavelet transform and wavelet coherence to geophysical time series. *Nonlinear Processes in Geophysics*, 11, 561-566.
4. Mokhov I.I., Bezverkhny V.A., Karpenko A.A., 2002: Mutual dynamics of atmospheric components and climatic characteristics during last 420,000 years from Vostok ice core. *Research Activity in Atmospheric and Oceanic Modelling*. H. Ritchie (ed.). WMO/TD-No.1105, 2.17-2.18.
5. Mokhov I.I., Bezverkhny V.A., Karpenko A.A., 2005: Diagnosis of relative variations in atmospheric greenhouse gas contents and temperature from Vostok Antarctic ice core paleoreconstructions. *Izvestiya, Atmospheric and Oceanic Physics*, 41(5), 523-592.
6. Mokhov I.I., Bezverkhny V.A., Karpenko A.A., 2010: Relative changes in temperature and concentration of greenhouse gases in the atmosphere from paleoreconstructions for last 800 ka. In: *Extreme Environmental Hazards and Catastrophes. V. I. Evaluation and Ways to Reduce Negative Consequences of Extreme Environmental Phenomena*. RAS, Moscow. 312-319. (in Russian)
7. Mokhov I.I., Bezverkhny V.A., Larkina I.S., 2015: Relative variations of temperature and aerosol concentration in the atmosphere from Antarctic ice cores. *Research Activities in Atmospheric and Oceanic Modelling*. E. Astakhova (ed.). WCRP Rep. No.12/2015, 2.4–2.5.