

Developing the next version of the Community Earth System Model: Aerosol effects using CMIP5 and CMIP6 forcing data.

*w/ Jean-François Lamarque, Andrew Gettelman, Richard Neale, Cecile Hannay,
and many others*

NCAR/CGD



Overview


- CESM2: New Physics
- Coupled model behavior with CMIP5 and CMIP6 emissions data

Evolution of CAM Model Physics

Model	CAM3 CCSM3	CAM4 CCSM4	CAM5 CESM1.0	CAM6 CESM2
Release	Jun 2004	Apr 2010	Jun 2010	Sometime 2017
PBL	Holtslag-Boville (1993)	Holtslag-Boville (1993)	Bretherton et al (2009)	CLUBB
Orographic form drag			Richter et al. (2010) “TMS”	Beljaars et al.2003
GW drag	McFarlane (1987)	McFarlane (1987)	McFarlane (1987) (non-orographic sources for WACCM)	Anisotropic/Low-level nonlinearities
Shallow Convection	Hack (1994)	Hack (1994)	Park et al. (2009)	CLUBB
Deep Convection	Zhang-McFarlane (1995)	Neale et al. (2008)	Neale et al. (2008)	Neale et al. (2008) Re-tuning
Microphysics	Rasch-Kristjansson (1998)	Rasch-Kristjansson (1998)	Morrison-Gottelman (2008)	Morrison-Gottelman v2 (2014)
Macrophysics	Rasch-Kristjansson (1998); Zhang et al (2003)	Rasch-Kristjansson (1998); Zhang et al (2003)	Park et al. (2011); Zhang et al (2003)	CLUBB
Radiation	Collins et al. (2001)	Collins et al. (2001)	Iacono et al. (2008)	Iacono et al. (2008)
Aerosols	Bulk Aerosol Model	Bulk Aerosol Model BAM	Modal Aerosol Model (MAM3, Ghan et al., 2011)	Modal Aerosol Model (MAM4, Liu et al., 2016)

 = switch parameterization

 = significant retuning/modification



***As of March 2017 development version
CESM2 looked pretty good***


- *Overall Taylor scores better than CESM1(LENS)*
- *Good El Niño*
- *Reasonable MJO*

As of March 2017 development version CESM2 looked pretty good

- *Overall Taylor scores better than CESM1(LENS)*
- *Good El Niño*
- *Reasonable MJO*

Next Steps:

- *Further development of land model*
- *Introduction of CMIP6 emissions*




What happened???

- Dust tuning and land model changes led to significantly colder model. ***(these changes are not considered optional)***
- CMIP6 emissions ????

Exp. 151 (Model ca. 5/2017;
emissions=CMIP5

Exp. 161 (Model ca. 5/2017;
emissions=CMIP6



What happened???


- CMIP6 emissions contribute, but may not be the whole story

RESTOM* Change in 20thC run with CMIP6 emissions

PD *PI*
161 (1951-1970) – (1851-1870)

OCEAN: CORR= 1.00; DEV=1.01; RMSE=2.56; BIAS=-0.06
LAND: CORR= 1.00; DEV=1.01; RMSE=3.52; BIAS=-0.93
GLOBAL: CORR= 1.00; DEV=1.01; RMSE=2.84; BIAS=-0.31

middle-top



ANN RESTOM

*RESTOM = Net SW – Net LW at “Top-of-model”


What else can we do???

- Patterns of RESTOM change suggestive of AIE
- ***2nd aerosol indirect effect (lifetime effect)***: more droplets → slower autoconversion → thicker longer-lived clouds
 - Poorly constrained. Some obs (e.g. in volcanic plumes) suggest is not very big

Using Volcanic Plume to study Aerosol-Cloud Interaction

CAM5.4 and CAM5.5 show big increases in cloud water with elevated aerosols. Here: an example from the Holuhraun eruption in Iceland in 2014.

Smaller drops are seen, but no increase in cloud water from Satellites (MODIS)



from A. Gettelman

Anomalies of Drop Size and Cloud Water for
October 2014 from long term mean.

Malavelle et al 2017, Nature

2nd indirect effect (thicker clouds from reduced autoconversion) seems to large in CAM

$$Au = kL^a N^b$$

Au =autoconversion rate

L =cloud liquid (kg/kg)

N =droplet number (#/kg)

Seifert&Beheng: $a=4$, $b=-2$

Khairouddinov&Kogan (KK); $a=2.67$, $b=-1.79$

- ***R. Wood***: b in KK could be as weak as -0.9.
- compromise $b=-1.1$.

Other tunable numbers (KK)


$k=0.01k_0$, $k=0.02k_0$ ($k_0=1350$)

“relative dispersion” can affect N entering Au calculation

Reducing droplet dependence in autoconversion improves simulation (w/resp HadCRUT)

Exp. 192 (Model ca. 9/2017;
emissions=CMIP5
 $Au = kL^a N^b$; with $b = -1.1$


Exp. 151 (Model ca. 5/2017;
emissions=CMIP5
Au=Seifert-Beheng



... but simulation forced with CMIP6 emissions
still doesn't warm enough (w/resp HadCRUT)


Exp. 192 (Model ca. 9/2017;
emissions=CMIP5
 $Au = kL^a N^b$; with $b = -1.1$)

Exp. 190 (Model ca. 9/2017;
emissions=CMIP6
 $Au = kL^a N^b$; with $b = -1.1$)



... but simulation forced with CMIP6 emissions
still doesn't warm enough (w/resp HadCRUT)


Exp. 161 (Model ca. 3/2017;
emissions=CMIP6)



... but simulation forced with CMIP6 emissions
still doesn't warm enough (w/resp HadCRUT)

Exp. 192 (Model ca. 9/2017;
emissions=CMIP5
 $Au = kL^a N^b$; with $b = -1.1$)

Exp. 190 (Model ca. 9/2017;
emissions=CMIP6
 $Au = kL^a N^b$; with $b = -1.1$)




Warming 1880-1920 to 1960-2000

192(CMIP5)

OCEAN: CORR= 1.00; DEV=0.98; RMSE=0.48; BIAS=0.33
 LAND: CORR= 1.00; DEV=0.99; RMSE=0.57; BIAS=0.32
 GLOBAL: CORR= 1.00; DEV=0.99; RMSE=0.52; BIAS=0.33

middle-top




ANN TS

Scale factor= 1.000

190(CMIP6)

OCEAN: CORR= 1.00; DEV=0.98; RMSE=0.40; BIAS=0.13
 LAND: CORR= 1.00; DEV=0.99; RMSE=0.47; BIAS=0.02
 GLOBAL: CORR= 1.00; DEV=0.99; RMSE=0.42; BIAS=0.10


middle-top



ANN TS


Scale factor= 1.000

192 minus 190



“Observed”

gisstemp Warming: (1880-1920) to (1960-2000)





SWCF



1900-1950

1950-2000



CMIP5



CMIP6



CMIP6 –
CMIP5



ANN SWCF


ANN SWCF

Scale factor= 1.000







Scale factor= 1.000

Differences in forcing CMIP5 -> CMIP6

Period=1970-1980



Approximate
location of cross-
sections on
following slide




Annual SWCF (W/m^2)

Annual Low Clouds (%)

Differences in forcing CMIP5 -> CMIP6

Longitude-height cross-sections along 26N (1970-1980 ann avg)




Differences in emissions

Surface SO_4 difference 1950-2000 CMIP6-CMIP5


OCEAN: CORR= 0.98; DEV=0.91; RMSE=10.87; BIAS=-1.44
LAND: CORR= 0.98; DEV=0.93; RMSE=25.34; BIAS=-2.28
GLOBAL: CORR= 0.98; DEV=0.94; RMSE=17.36; BIAS=-1.84

middle-top




ANN so4_a1_SRF

Scale factor= 1.000e+11



from J.F.Lamarque


Transport of sulfate from US SO₂ emissions




SWCF CMIP6-CMIP5

Conley et al., submitted, 2017


“Accelerated development” runs






B1850

BHIST



19xx = 1930, 1950 ...




19xx = 1930, 1950 ...

1950 Starts vs 1850 Starts


CMIP5 emiss.

Annual mean TS Global



CMIP6 emiss.


Annual mean TS Global



1930 Starts vs 1850 Starts


Latest (last week) model w/
CMIP6 emiss.

Annual mean TS Global



CMIP5 emiss.

Annual mean TS Global





197_1930

197



192_1930

192

1940-1960




1960-1980



difference

$r(\text{diff}) = 0.49$




ANN TS
Scale factor= 1.000

ANN TS
Scale factor= 1.000

$r()=0.07$
 $r()=0.12$

$r(\text{diff}) = 0.72$



ANN TS
Scale factor= 1.000

ANN TS
Scale factor= 1.000



197_1930

197



192_1930

192

1940-1960




1960-1980



difference


$r(\text{diff}) = 0.39$



$r()=0.13$

$r()=0.05$

$r(\text{diff}) = 0.48$



ANN SWCF

ANN SWCF

ANN SWCF

ANN SWCF

Scale factor= 1,000

Scale factor= 1,000

Scale factor= 1,000

Scale factor= 1,000

Possibilities

- Model is too sensitive to aerosol details
- Problem with CMIP5 (and CESM2) or CMIP6 emissions

Summary and Plans

- Current CESM2 performs well when forced with CMIP5 emission data, not so with CMIP6 emissions
- East Pacific feedbacks may play a role in cooling later 20th Century in CMIP6 - mechanism unclear.
- Beginning more intensive assessment of individual CMIP6 aerosol components
- Beginning physics “swaps”, e.g., PBL, ShCu schemes
- Examining changes to aerosol activation parameterizations

Thank You

Overview of CLUBB's solution procedure

Advance 10 prognostic equations

$$\overline{w}, \overline{\theta_l}, \overline{q_t}, \overline{w'^2}, \overline{w'^3}, \overline{q_t'^2}, \overline{\theta_l'^2}, \overline{q_t'\theta_l'}, \overline{w'q_t'}, \overline{w'\theta_l'}$$

Note:

No equations for

$$\overline{u'w'}$$

$$\overline{v'w'}$$

Close dissipation terms

Δt

Use PDF to close higher-order moments, buoyancy terms

$$\overline{w'q_t'^2}, \overline{w'\theta_l'^2}, \overline{w'q_t'\theta_l'}, \overline{w'^2q_t'}, \overline{w'^2\theta_l'}, \overline{w'^4},$$

$$\overline{q_t'\theta_v'}, \overline{\theta_l'\theta_v'}, \overline{w'\theta_v'}, \overline{w'^2\theta_v'}$$

Select PDF from given functional form to match 10 moments



Bi-normal PDFs



Diagnose cloud fraction, liquid water from PDF

Annual mean shortwave cloud forcing

CAM5.4 (before CLUBB)



f.e15b03wch.F55W5SC.f09_f09.C54-OGW-T1-x03 (yrs 1980-1989)

TOA SW cloud forcing mean = -50.06 W/m^2





CERES-EBAF

TOA SW cloud forcing mean = -47.15 W/m^2



f.e15b03wch.F55W5SC.f09_f09.C54-OGW-T1-x03 - CERES-EBAF



mean = -2.91 rmse = 12.83 W/m^2



CAM5.5 (uses CLUBB)



f.e15b03wch.F55W5SC.f09_f09.CLB-OGW-T1-x01 (yrs 1980-1989)

TOA SW cloud forcing mean = -48.59 W/m^2





CERES-EBAF

TOA SW cloud forcing mean = -47.15 W/m^2



f.e15b03wch.F55W5SC.f09_f09.CLB-OGW-T1-x01 - CERES-EBAF



mean = -1.44 rmse = 10.22 W/m^2



Anisotropic Orographic Gravity Wave Drag (scales~>10km)

Ridge orientation determines wave orientation and direction of drag force not low-level wind.

Ridge height estimate is based on min and max elevations of mean ridge profile not based on subgrid variance




Generation of ancillary topography files

- No traceable process for generating topography forcing data existed for CESM1 or earlier versions.
 - Derivation of subgrid variables and smoothing of mean elevations left up to dycore developers. ***Note: all dycores employ additional smoothing beyond binning to grid.***
- New procedure starts from 1km GMTED2010 data (or GTOPO30) mapped to 3km cubed sphere grid. Further processing follows from 3km cubed sphere topo (*Lauritzen et al. GMD 2015*)

Temperature (Annual) - AMIP

- *Lower stratospheric 'cold-pole' problems see significant decrease*
- Polar near-surface, improved stable PBLs
- Tropics go warm, cold, warm, cold, colder, cold
- A response to continued changes to deep convection
- Trade-off for climate mean and variability



Slide from Rich Neale


MG2 and MAM4

- Prognostic precipitation species (*rain and snow*)
- Additional carbon aerosol mode with longer residence time (*increases transport to high-latitudes*)

Coupled system (CESM)


Description of progress through ~March 2017 (***configuration 125***)

Skill Score (current simulation:#125)





- General monotonic improvement from CESM1 (DJF/ANN)
- Large initial degradation in JJA mostly recovered
- Removing super-saturation -> improved skill, but high climate sensitivity
- Land model strongly impacts JJA score (new land at 118).

ENSO in CESM2



Tropical Variability (Precipitation) - CESM




- Largest variation in MJO power
- Increased power and zonal wavenumber extent

Madden Julian Oscillation (MJO)

Precipitation

850-mb U




Precipitation

- Lag correlation with Indian-Ocean precip
- 20-100day band pass filter, 10S-10N
- 9 years, DJFMAM


OLR

Differences in emissions






Bootstrap analysis of ΔSWCF





10000 artificial timeseries generated from
20 year 195.001 and 195.011(CMIP5).
Random sampling w/ replacement: 3yr, 5yr,
10yr and 16yr artificial timeseries




CMIP5 (mean=-1.04)

CMIP6 (mean=-1.39)






OCEAN: CORR= 1.00; DEV=1.00; RMSE=1.68; BIAS=-0.05
LAND: CORR= 1.00; DEV=1.01; RMSE=1.48; BIAS=-0.41
GLOBAL: CORR= 1.00; DEV=1.00; RMSE=1.63; BIAS=-0.16
middle-top



ANN SWCF

Scale factor= 1.000

OCEAN: CORR= 1.00; DEV=1.00; RMSE=2.06; BIAS=-0.57
LAND: CORR= 1.00; DEV=1.01; RMSE=1.81; BIAS=-0.85
GLOBAL: CORR= 1.00; DEV=1.00; RMSE=1.99; BIAS=-0.65
middle-top



ANN SWCF

Scale factor= 1.000

PBL Form Drag (from smooth small obstacles)

TMS:

$$\mathbf{F}_x = C_D |\mathbf{U}| U(z_{LM}), \quad C_D = k_\delta \ln \left(\frac{z_{LM}}{z_0} \right) \frac{\rho U^2}{2}, \quad z_0 \propto \sqrt{\langle h_\delta^2 \rangle}$$





- Logarithmic in h'_δ . Only applied in lowest model layer.

Beljaars et al. (2004):

$$\mathbf{F}_x = -\alpha \beta C_{md} C_{corr} |\vec{U}(z)| \vec{U}(z) 2.109 e^{-(z/1500)^{1.5}} a_2 z^{-1.2}, \quad a_2 \propto \langle h_\delta^2 \rangle$$


- Proportional to $h'_\delta{}^2$. Applied over physically based vertical profile.

h_δ are topographic perturbations with scales below 3km derived from GMTED data. $\langle \rangle$ represents averaging to model grid.

ERA-I**ERA-I****IOGW****AOGW**


Taylor Diags XTR : CORR= 0.97; DEV=1.00; RMSE=2.81; BIAS=0.94
Taylor Diags SH : CORR= 0.97; DEV=1.01; RMSE=3.19; BIAS=1.80
Taylor Diags NH : CORR= 0.98; DEV=1.08; RMSE=2.38; BIAS=0.07

Middle - Top




Taylor Diags XTR : CORR= 0.98; DEV=0.96; RMSE=2.52; BIAS=0.35
Taylor Diags SH : CORR= 0.98; DEV=1.01; RMSE=2.77; BIAS=1.43
Taylor Diags NH : CORR= 0.98; DEV=0.99; RMSE=2.25; BIAS=-0.72


Middle - Top

**Zonal-mean zonal wind (DJF)**


ERA-I




ERA-I




IOGW




AOGW



Taylor Diags XTR : CORR= 0.98; DEV=1.16; RMSE=2.87; BIAS=0.71
Taylor Diags SH : CORR= 0.99; DEV=1.17; RMSE=3.80; BIAS=2.20
Taylor Diags NH : CORR= 0.99; DEV=0.97; RMSE=1.42; BIAS=-0.77



Taylor Diags XTR : CORR= 0.99; DEV=1.09; RMSE=2.09; BIAS=0.14
Taylor Diags SH : CORR= 0.99; DEV=1.10; RMSE=2.33; BIAS=1.35
Taylor Diags NH : CORR= 0.98; DEV=0.95; RMSE=1.82; BIAS=-1.08



Zonal-mean zonal wind (SON)

Short cuts ???


- Full 20th century coupled runs long time: ~8-10 days to go from 1850 to 2005

Climatological year 2000 runs

20 year runs with PD (year 2000) and
PI(year 1850) aerosols


All other forcings, e.g. GHG, SSTs etc.
are for year 2000

Change in SWCF from PI to PD



Mean change in SWCF from PI to PD

CMIP5 (mean=-1.04)



CMIP6 (mean=-1.39)

